

International Journal of Scientific Research and Reviews

A Review on the Synthesis and Use of Benzobisthiazoles: An Important Class of Heterocycles

Ratna Mukherjee *

Department of Chemistry, Dum Dum Motijheel College, 1, Motijheel Avenue, Kolkata 700074,
West Bengal, India

Email: ratna.org@gmail.com

ABSTRACT

This review emphasises on the synthesis of various benzobisthiazole derivatives both linear and angular isomers. According to the position of nitrogen and sulphur atoms in this type of heteroarene, the linear benzobisthiazole moiety has two types of skeleton whereas the angular isomer has three types. Additionally, this review enlightens the role of few benzobisthiazole derivatives in biological field. Eventually a wide variety of other uses of benzobisthiazoles and their polymers are also discussed in brief.

KEYWORDS: benzobisthiazoles, synthesis, biological activity

***Corresponding author**

Dr. Ratna Mukherjee

Assistant Professor, Department of Chemistry

Dum Dum Motijheel College

1, Motijheel Avenue, Kolkata-700074, West Bengal, India

E-mail: ratna.org@gmail.com

1. INTRODUCTION

The importance of sulphur atoms in the living organisms is well documented.¹ The thiazoles constitute an important class of heterocyclic that contains one sulphur atom. The thiazoles, both substituted and annulated, are present in many natural and synthetic products possessing a wide range of pharmacological activities. These molecules display, *inter alia*, antiviral, anti-ulcer, antibacterial, antifungal, anticonvulsant, antiparkinsonian and anti-inflammatory activities. Some of them are vitamin B₁,^{2,3} and the bioactive natural products are camalexins (antimicrobial) and spiobrassinins (antifungal).⁴

Some examples of thiazole containing synthetic drugs are the anticonvulsant riluzole, the antischistosomal niridazole, the anthelmintic thiabendazole, the antimicrobial sulfathiazole and the anti-HIV ritonavir.² The well known synthetic drugs famotidine and nizatidine (anti-ulcer)⁵ also deserve special mention. Some of the synthetic thiazolyliindoles also show impressive, selective COX-2 (cyclooxygenase) inhibitory activity.^{6,7} The antiparkinsonian talipexole is also a well known bioactive compound.²

Thiazole and thiazole-annulated scaffolds are among the most prominent building blocks in chromophores with potential applications in fields such as nonlinear optics (NLO),⁸⁻¹⁶ organic light-emitting diodes,¹⁷ organic field-effect transistors,¹⁸⁻²⁰ dye-sensitized solar cells,^{21,22} polymers,²³ liquid crystals,²⁴ photonucleases,²⁵ fluorescent dyes,^{26,27} insecticides²⁸ and antioxidants.²⁹

Among the fused thiazoles, benzothiazoles are of considerable interest as a result of their important biological and biophysical (nonlinear optical materials, molecular dyads, chemosensors)^{30,31} properties. 2-Aryl and 2-heteroaryl substituted benzothiazoles have been studied as antitumor,³² antimicrobial³³ and antifungal agents³⁴ and as imaging agents for β -amyloid.³⁵ Pertinently, 2-(4-aminophenyl)benzothiazoles are fast emerging as promising antitumor drugs.³⁶⁻³⁸



Among the annulated thiazoles, benzobisthiazoles have been less researched; a few reports have indicated that derivatives of these compounds show some potent biological activity. So in the present review benzobisthiazole was chosen in order to discuss about their synthesis, biological activity and other miscellaneous uses.

2. EXTANT SYNTHESSES OF BENZOBISTHIAZOLES

Two types of linearly fused and three types of angularly fused benzobisthiazoles can, in principle, be formed by the fusion of the thiazole and the benzothiazole nuclei. These parent tricycle molecules are.

Linear benzobisthiazoles:

1) Benzo[1,2-*d*:4,5-*d'*]bisthiazoles (Type I) 2) Benzo[1,2-*d*:5,4-*d'*]bisthiazoles (Type II)

Angular benzobisthiazoles:3) Benzo[1,2-*d*:3,4-*d'*]bisthiazoles (Type III) 4) Benzo[1,2-*d*:4,3-*d'*]bisthiazoles (Type IV)5) Benzo[1,2-*d*:6,5-*d'*]bisthiazoles (Type V)**Fig. 1. Linear Benzobisthiazoles****Fig. 2. Angular Benzobisthiazoles**

All the five classes of benzobisthiazoles are in fact reported in the literature. In the majority of the reported syntheses of these compounds, either *ortho*-, *meta*- and *para*-phenylenediamines or substituted 5/6/7-aminobenzothiazoles were the starting materials. The reported syntheses of the different classes of the benzobisthiazoles (Types I-V) are briefly discussed in this review.


2.1 Preparation of Benzo[1,2-*d*:4,5-*d'*]bisthiazoles (Type I)

In 1903, Green and Perkin³⁹ first claimed to have prepared this class of linear benzobisthiazoles. Later, however, three different groups⁴⁰⁻⁴² proved that the benzobisthiazoles prepared by Green and Perkin did indeed belong to the angular [1,2-*d*:4,3-*d'*] series, i.e. Type IV. These latter workers based their conclusions on UV, IR and NMR spectroscopic analyses.

The benzobisthiazoles of Type I were prepared from *p*-phenylenebisthiourea⁴⁰ (**Scheme 1**) and 5-amino-6-nitrobenzothiazole⁴¹ (**Scheme 2**). Type I benzobisthiazoles were also prepared by derivatisations of other benzobisthiazoles.⁴³ Some syntheses of Type I benzobisthiazoles are discussed here.

2.1.1 Starting from *p*-Phenylenebisthiourea⁴⁰


Bromine-catalysed cyclisation of thioureas derived from aromatic amines is one of the oldest (1903) methods, known as Hugershoff reaction.⁴⁴ This method was again brought to fore by Ambati et al. in 1997.⁴⁵ Application of Hughschhof protocotol to the synthesis of 2,6-diaminobenzo[1,2-*d*:4,5-*d'*]bisthiazole starting from *p*-phenylene-bisthiourea is depicted in **Scheme 1**.⁴⁰



Scheme 1

2.1.2 Starting from 5-Amino-6-nitrobenzothiazole⁴¹

5-Amino-6-nitrobenzothiazole was sequentially diazotized and treated with sodium disulfide, the resulting disulfide reductively cleaved to the corresponding aminothiophenol and finally cyclised to form 2,6-dimethylbenzo[1,2-*d*:4,5-*d'*]bisthiazole (**Scheme 2**).



Scheme 2

2.2 Preparation of Benzo[1,2-*d*:5,4-*d'*]bisthiazoles (Type II)


In 1922, S.R.H. Edge⁴⁶⁻⁴⁸ claimed to have prepared 2,6-dimethyl[1,2-*d*:5,4-*d'*]bisthiazole, i.e. a member of Type II. Subsequently, it was demonstrated^{49,50} that the product was indeed the angular isomer, viz. 2,7-dimethylbenzo[1,2-*d*:3,4-*d'*]bisthiazole (Type III).

Various members of Type II benzobisthiazoles were synthesised from different precursors, viz. 1,3-phenylenediamine,^{43,40} 2,5-diaminobenzothiazole,⁴⁰ 1,3-dinitro-4,6-dithiocyanobenzene,⁴⁰

dinitrothioresorcinol,⁴⁹ *m*-phenylenebisthiourea⁵¹ and 1,3-dichloro-4,6-dinitrobenzene.^{50,52} Many of these products were further derivatised, leading to the formation of a number of other members of this Type.⁴³ Some of these syntheses are shown in this review.

2.2.1 Starting from *m*-Phenylenediamine^{43,40}

m-Phenylenediamine was made to undergo successive thiocyanation and cyclisation, as depicted below, to result in the formation of 2,6-diaminobenzo[1,2-*d*:5,4-*d'*]bisthiazole (**Scheme 3**).




Scheme 3

The corresponding bisdiethylaminoacetyl derivative was also prepared from this benzobisthiazole.⁴³

2.2.2 Starting from 2,5-Diaminobenzothiazole⁴⁰


Following the same protocol as above but using 2,5-diaminobenzothiazole as the substrate, 2,6-diaminobenzo[1,2-*d*:5,4-*d'*]bisthiazole was prepared (**Scheme 4**).



Scheme 4

2.2.3 Starting from 1,3-Dinitro-4,6-dithiocyanobenzene⁴⁰


Preformed 1,3-Dinitro-4,6-dithiocyanobenzene was subjected to reductive cyclisation to furnish 2,6-diaminobenzo[1,2-*d*:5,4-*d'*]bisthiazole in a low yield (**Scheme 5**).⁴⁰ The low yield of the product renders this method of only academic interest.



Scheme 5

2.2.4 Starting from Dinitrothioresorcinol⁴⁹


2,4-Dinitrothioresorcinol was reduced and acetylated to furnish the 2,6-dimethylbenzo[1,2-*d*:5,4-*d'*]bisthiazole in a moderate yield (**Scheme 6**).



Scheme 6

2.2.5 Starting from *m*-Phenylenebisthiourea⁵¹


m-Phenylenediamine (*m*-PD) dihydrochloride on thiocyanation furnished the bisthiourea which was cyclised by Hughsorhoff reaction to furnish 2,6-diamino-benzo[1,2-*d*:5,4-*d'*]bisthiazole (**Scheme 7**).



Scheme 7

2.2.6 Starting from 1,3-Dichloro-4,6-dinitrobenzene^{50,52}

Various members of Type II were prepared using 1,3-dichloro-4,6-dinitro-benzene as the starting material using different methodologies.^{50,52} Only one example is depicted in **Scheme 8**. Here 2,6-dimethylbenzo[1,2-*d*:5,4-*d'*]bisthiazole was prepared (**Scheme 8**).⁵⁰




Scheme 8

a. Preparation of Benzo[1,2-*d*:3,4-*d'*]bisthiazoles (Type III)

2.3.1 Starting from *m*-Phenylene Di Thioacetamide


An example of this type of benzobisthiazoles was first reported to be prepared by Edge⁴⁶⁻⁴⁸ by the oxidative ring closure of *m*-phenylenedithioacetamide employing Jacobson reaction.⁵³ The latter is the oxidative cyclisation of aryl thioamides by aqueous alkaline potassium ferricyanide to form thiazoloarenes. But Edge wrongly identified the product as the linear molecule, 2,6-dimethylbenzo[1,2-*d*:5,4-*d'*]bisthiazole (Type II). Later G. Grandolini et al.⁴⁹ and O. Neunhoeffter et al.⁵⁰ prepared the aforesaid molecule by two different methods (**Schemes 6** and **8**, respectively), already discussed before, and found their melting points to be different from that of Edge's compound. They, therefore, opined that the compound prepared by Edge did indeed bear the only possible alternative angular structure, viz. 2,7-dimethylbenzo[1,2-*d*:3,4-*d'*]bisthiazole (Type III). The actual experiment undertaken by Edge is represented in **Scheme 9**.



Scheme 9

2.3.2 Starting from 5/7-Aminobenzothiazoles⁵⁴⁻⁵⁶

Various benzobisthiazoles of Type III were prepared from 5/7-amino- benzothiazoles.⁵⁴⁻⁵⁶ A particular example is depicted in **Scheme 10**. Here 2-amino-7-methylbenzo[1,2-*d*:3,4-*d'*] bisthiazole and 2-amino-5-methylbenzo[1,2-*d*:3,4-*d'*] bisthiazole, were prepared in one-pot reaction.⁵⁵




(i): $\text{NH}_4\text{SCN}/\text{AcOH}$, $\text{O}=\text{C}(\text{NHCl})_2 / \text{AcOH}$, ice-cooled, 1-2 h

Scheme 10

2.3.3 Starting from 2-Methyl(benzothiazol-5-yl)thiourea⁵⁷

2-Methyl-5-thioureidobenzothiazole was cyclised by Hugershoff reaction to furnish 2-amino-7-methylbenzo[1,2-*d*:3,4-*d'*]bisthiazole (**Scheme 11**).




Scheme 11

2.4 Preparation of Benzo[1,2-*d*:4,3-*d'*]bisthiazoles (Type IV)

2.4.1 Starting from *p*-Phenylenediamine


Green and Perkin³⁹ were the first to claim to have achieved the synthesis of 2,6-disubstituted benzo[1,2-*d*:4,5-*d'*]bisthiazoles (Type I) starting from *p*-phenylenediamine (*p*-PD). They thionated *p*-PD and assumed that the product was 1,4-diaminobenzene-2,5-dithiosulfonic acid (**I**). In other words, they assumed that in the dithionated product, (i) each thiosulfonic group was *ortho*- to an amino group and (ii) the two thiosulfonic acid groups were *para*- to each other and hence believed that the resulting cyclised products were the aforesaid linear benzobisthiazoles, **IIa** and **IIb** (Type I). The reactions are depicted in **Scheme 12**.



Scheme 12

In the late sixties, three independent groups⁴⁰⁻⁴² synthesised 2,6-dimethyl, 2,6-diamino- and 2,6-dimethylaminobenzo[1,2-*d*:4,5-*d'*]bisthiazoles following established literature procedures. They also repeated Green and Perkin's work. From a consideration of the UV, IR and NMR spectra as well as the dipole moments of both sets of benzobisthiazoles, they demonstrated that the compounds prepared by Green and Perkin were actually the angular products, i.e. 2,7-disubstituted benzo[1,2-*d*:4,3-*d'*]bisthiazoles (Type IV).

It thus became evident from the aforesaid studies that the presumed 1,2,4,5-configuration of the intermediate diaminobenzene dithiosulfonic acid (**I**) was wrong. This product had indeed the 1,2,3,4-tetrasubstituted benzene structure (**III**), and the resulting products were naturally the angular isomers (**IVa,b**). The actual work undertaken by Green and Perkin is represented in **Scheme 13**.




Scheme 13

It requires to be brought to the fore that, subsequently to Green and Perkin's work,³⁹ some reports^{58,59} came out where the authors prepared benzobisthiazoles starting from diaminobenzene dithiosulfonic acid, following the report of Green and Perkin.³⁹ Now that the structure of the disulfonic acid was revised, the benzobisthiazoles prepared by these workers did not have the linear structures (Type I) claimed by them; these were actually the angular benzobisthiazoles (Type IV).

2.4.2 Staring from 6-Aminobenzothiazole⁶⁰


Two benzobisthiazoles of Type IV were prepared from 6-amino and 2-methyl-6-aminobenzothiazoles by thiocyanation (using ammonium thiocyanate, acetic acid and dichlorourea), followed by acid-catalysed cyclisation (**Scheme 14**).⁶⁰



Scheme 14

2.5 Preparation of Benzo[1,2-d:6,5-d']bisthiazoles (Type V)⁴⁰

A member of this class of benzobisthiazoles was prepared by the cyclisation of *ortho*-phenylenebis(*N'*-methylthiourea) by cyclisation with bromine or sulphuryl chloride in chloroform (**Scheme 15**).



Scheme 15

Moreover, other members of Type V were also prepared by derivatisation of members of this class.⁴³


Following the above literature in 2010 Chakrabarty et al.⁶¹ have reported a regioselective synthesis of various 2-substituted benzobisthiazoles by more or less a unified approach starting from 6-amino-benzothiazole. The amine was converted to: (i) 2-aminobenzobisthiazole by direct cyclisation (ii) 2-alkylamino/anilinobenzobisthiazoles via the corresponding 6-(*N'*-alkyl/phenyl)thioureidobenzothiazoles (iii) 2-alkylthio-benzobisthiazoles via the intermediacy of alkyl (benzothiazol-6-yl)dithiocarbamates and (iv) 2-alkylbenzobisthiazoles from the *N*-(6-benzothiazolyl) thioamides.

In the same year Mike et al.⁶² have reported the synthesis of several synthetically useful 2,6-disubstituted benzobisthiazoles using Lewis acid as catalyst. It was a ring-closing reaction between substituted orthoesters and diamino benzene dithiol. The resulting benzobisthiazoles were obtained in good yields.

3. BIOLOGICAL ACTIVITY

A few references revealed that the benzobisthiazoles derivatives demonstrate potent anti-inflammatory, antibacterial, antifungal and antiprotozoal activity.^{59,43,63} In general, linear benzobisthiazoles have been found to display higher biological activity than their angular counterparts.^{59,43}

In 1992, Cullen et al.⁴³ have reported that 2,6-bis(*N,N*-diethylamino) benzo [1,2-*d*:5,4-*d'*] bisthiazole was found to inhibit the swelling of the uninjected paw in the prophylactic adjuvant arthritis model with an ED₅₀ of 2.3 mg/kg orally.



Few years later Chakrabarty et al.⁶¹ have investigated the anti-inflammatory potential of benzobisthiazole derivatives by measuring their ability to inhibit cyclooxygenase (COX). The

benzobisthiazoles were screened for inhibition of both COX-1 and COX-2 using naproxen⁶⁴ as the standard following a recent protocol.^{65,66} In the benzobisthiazole series, the methylamino derivative was most active, the anilino and the methylthio derivatives were less active and the rest displayed poor activity. 2-Methylthiobenzobisthiazole exhibited the best COX-2 inhibition (36%), followed by the 2-amino derivative. Although the 2-substituted benzobisthiazoles possessed the lowest COX-1 inhibitory activity, they displayed better COX-2 inhibition. All the synthesised compounds are less potent COX-1 and COX-2 inhibitors than naproxen.

In 2013 in another report⁶⁷, angular benzobisthiazole derivatives with a benzamide substitution (phenyl ring bearing different substitution) were recognized as novel helicase inhibitors through high throughput screening against purified *Staphylococcus aureus* and *Bacillus anthracis* replicative helicases. Initial investigation of structure-activity relationships (SARs) focused on the benzamide portion of the benzobisthiazole scaffold.

In 2016, again angular benzobisthiazole moiety was identified as selective and potent inhibitor for CLK protein kinase family members by Prak et al.⁶⁸

4. MISCELLANEOUS APPLICATIONS

- In 2015, it was reported that organic benzobisthiazolium⁶⁹ salts bring various advantages in the field of Non Linear Optical materials, due to their improved solubility, greater stability (including photostability) and mainly enhanced Intramolecular Charge Transfer.
- In 2009 Ahmed et al.⁷⁰ recognized that the incorporation of benzobisthiazole into a regioregular poly(3-alkylthiophene) improve the oxidative stability by increasing the ionization potential (IP), thermal stability, inter chain interactions and thus enhance the charge transport properties of the polymers. The new copolymer semiconductor indeed showed an increased ionization potential and robust thermal and air stability.
- Polybenzobisthiazole were used in electronic applications as electron transport materials⁷¹ in organic light-emitting diodes,⁷² as nonlinear optical materials,⁷³⁻⁷⁵ and as charge photogeneration materials in xerographic photoreceptors.⁷⁶⁻⁷⁸ Light-emitting diodes containing polybenzobisthiazole electron transport layer were found to be stable in air for over a period of 10 months.⁷¹
- Materials based on benzobisthiazole units also showed other attractive electronic properties, such as high fluorescence, thermal stability, high electron affinity, electroluminescence properties,^{79,80} and nonlinear optical properties.^{72,81-89} As a result of these properties, such

materials have been used as electron transport materials in OLEDs⁷², charge photogeneration materials^{90,77} field-effect transistors and solar cells.⁹¹⁻⁹³

5. CONCLUSION

This review article provides an extensive overview for the synthesis of various substituted benzobisthiazoles. Few biological applications of benzobisthiazoles and its derivatives were also brought to vanguard. Besides different uses of benzobisthiazole moiety as well as their polymers was discussed in brief. The current review is significant and holds the promise of being useful in the study of benzobisthiazoles.

6. REFERENCES

1. Fontecave M, Ollagnier-de-Choudens S, Mulliez E. Biological Radical Sulfur Insertion Reactions. *Chem. Rev.* 2003; 103(6): 2149-2166.
2. Kleemann A. Pharmaceutical substances. 4th edn. George-Thieme: New York; 2001.
3. De Souza MVN. Synthesis and biological activity of natural thiazoles : an important class of heterocyclic compounds. *J. Sulfur Chem.* 2005; 26(4-5): 429-449.
4. Pedras MSC, Okanga FI, Zaharia IL et al. Phytoalexins from crucifers: synthesis, biosynthesis, and biotransformation. *Phytochemistry.* 2000; 53(2): 161-176.
5. Akita H, Sutou N, Sasaki T et al. Alternative synthesis of cystothiazole A. *Tetrahedron.* 2006; 62(50): 11592-11598.
6. Faucher AM, Bailey MD, Beaulieu PL et al. Synthesis of BILN 2061, an HCV NS3 Protease Inhibitor with Proven Antiviral Effect in Humans. *Org. Lett.* 2004; 6(17): 2901-2904.
7. Woods W, Mc Croskey RW, Michaelides MR et al. Thiazole analogues of the NSAID indomethacin as selective COX-2 Inhibitors. *Bioorg. Med. Chem. Lett.* 2001; 11(10): 1325-1328.
8. Breitung EM, Shu CF, Mc Mahon RJ. Thiazole and Thiophene Analogues of Donor-Acceptor Stilbenes: Molecular Hyperpolarizabilities and Structure-Property Relationships. *J. Am. Chem. Soc.* 2000; 122(6): 1154-1160.
9. Hrobárik P, Zahradník P, Fabian WMF. Computational design of benzothiazole-derived push-pull dyes with high molecular quadratic hyperpolarizabilities. *Phys. Chem. Chem. Phys.* 2004; 6(3): 495-502.
10. Hrobárik P, Sigmundová I, Zahradník P et al. Molecular Engineering of Benzothiazolium Salts with Large Quadratic Hyperpolarizabilities: Can Auxiliary Electron-Withdrawing Groups Enhance Nonlinear Optical Responses?. *J. Phys. Chem. C.* 2010; 114(50): 22289-22302.

11. Ma XH, Ma F, Zhao ZH et al. Toward highly efficient NLO chromophores: Synthesis and properties of heterocycle-based electronically gradient dipolar NLO chromophores. *J. Mater. Chem.* 2010; 20(12): 2369-2380.
12. Raposo MMM, Castro MCR, Belsley M et al. Push-pull bithiophene azo-chromophores bearing thiazole and benzothiazole acceptor moieties: Synthesis and evaluation of their redox and nonlinear optical properties. *Dyes Pigm.* 2011; 91(3): 454-465.
13. El-Shishtawy RM, Borbone F, Al-Amshany ZM et al. Thiazole azo dyes with lateral donor branch: Synthesis, structure and second order NLO properties. *Dyes Pigm.* 2013; 96(1): 45-51.
14. Hrobáriková V, Hrobárik P, Gajdoš P et al. Benzothiazole-Based Fluorophores of Donor- π -Acceptor- π -Donor Type Displaying High Two-Photon Absorption. *J. Org. Chem.* 2010; 75(9): 3053-3068.
15. Hrobárik P, Hrobáriková V, Sigmundová I et al. Benzothiazoles with Tunable Electron-Withdrawing Strength and Reverse Polarity: A Route to Triphenylamine-Based Chromophores with Enhanced Two-Photon Absorption. *J. Org. Chem.* 2011; 76(21): 8726-8736.
16. Belfield KD, Morales AR, Kang BS et al. Synthesis, Characterization, and Optical Properties of New Two-Photon-Absorbing Fluorene Derivatives. *J. Chem. Mater.* 2004; 16(23): 4634-4641.
17. Yu G, Yin SW, Liu YQ et al. Structures, Electronic States, and Electroluminescent Properties of a Zinc(II) 2-(2-Hydroxyphenyl) benzothiazolate Complex. *J. Am. Chem. Soc.* 2003; 125(48): 14816-14824.
18. Ando S, Murakami R, Nishida J et al. n-Type Organic Field-Effect Transistors with Very High Electron Mobility Based on Thiazole Oligomers with Trifluoromethylphenyl Groups. *J. Am. Chem. Soc.* 2005; 127(43): 14996-14997.
19. McEntee GJ, Vilela F, Skabara PJ et al. Self-assembly and charge transport properties of a benzobisthiazole end-capped with dihexyl thienothiophene units. *J. Mater. Chem.* 2011; 21(7): 2091-2097.
20. Lin Y, Fan H, Li Y et al. Thiazole-based organic semiconductors for organic electronics. *Adv. Mater.* 2012; 24(23): 3087-3106.
21. Zhang W, Feng Q, Wang ZS et al. Novel thiazolo [5,4-d] thiazole-based organic dyes for quasi-solid-state dye-sensitized solar cells. *Chem.-Asian J.* 2013; 8(5): 939-946.

22. Saeki A, Tsuji M, Yoshikawa S et al. Boosting photovoltaic performance of a benzobisthiazole based copolymer: a device approach using a zinc oxide electron transport layer. *J. Mater. Chem. A*. 2014; 2(17): 6075-6080.
23. Wang LY, Zhang CX, Liu ZQ et al. A 2-D ladder-type polymer, $\text{Mn}_2(\text{NIT2-thz})_2\text{Cl}_4(\text{H}_2\text{O})_2$: synthesis, crystal structure and magnetic properties. *Inorg. Chem. Commun.* 2003; 6(9): 1255-1258.
24. Al-Dujali AH, Atto AT, Atkurde AM. Synthesis and liquid crystalline properties of models and polymers containing thiazolo [5,4-d] thiazole and siloxane flexible spacers. *Eur. Polym. J.* 2001; 37(5): 927-932.
25. Li Y, Xu Y, Qian X, Qu B. Naphthalimide-thiazoles as novel photonucleases: molecular design, synthesis, and evaluation. *Tetrahedron Lett.* 2004; 45(6): 1247-1251.
26. Tintcheva I, Maximova V, Deligeorgiev T et al. New asymmetric monomethine cyanine dyes for nucleic-acid labelling: absorption and fluorescence spectral characteristics. *J. Photo hem. Photobiol. A: Chem.* 2000; 130(1): 7-11.
27. Rueker VC, Foister S, Melander C, Dervan PB. Sequence Specific Fluorescence Detection of Double Strand DNA. *J. Am. Chem. Soc.* 2003; 125(5): 1195-1202.
28. Wang Q, Li H, Li Y, Huang R. Synthesis and Herbicidal Activity of 2-Cyano-3-(2-chlorothiazol-5-yl) methylaminoacrylates. *J. Agric. Food. Chem.* 2004; 52(7): 1918-1922.
29. Yanagimoto K, Lee KG, Ochi H, Shibamoto T. Antioxidative Activity of Heterocyclic Compounds Found in Coffee Volatiles Produced by Maillard Reaction. *J. Agric. Food. Chem.* 2002; 50(19): 5480-5484.
30. Batista RMF, Costa SPG, Malheiro EL, Belsley M, Raposo MMM. Synthesis and characterization of new thienylpyrrolyl-benzothiazoles as efficient and thermally stable nonlinear optical chromophores. *Tetrahedron.* 2007; 63(20): 4258-4265.
31. Reddy PVG, Lin YW, Chang H-T. Synthesis of novel benzothiazole compounds with an extended conjugated system. *ARKIVOC.* 2007; (xvi): 113-122.
32. Mortimer CG, Wells G, Crochard JP et al. Antitumor Benzothiazoles. 26.¹ 2-(3,4-Dimethoxyphenyl)-5-fluorobenzothiazole (GW 610, NSC 721648), a Simple Fluorinated 2-Arylbenzothiazole, Shows Potent and Selective Inhibitory Activity against Lung, Colon, and Breast Cancer Cell Lines. *J. Med. Chem.* 2006; 49(1): 179-185.
33. Akbay A, Oren I, Temiz-Arpaci O, Aki-Sener E, Yalcin I. Synthesis and HIV-1 reverse transcriptase inhibitor activity of some 2, 5, 6-substituted benzoxazole, benzimidazole, benzothiazole and oxazolo(4,5-b)pyridine derivatives. *Arzneim.Forsch./Drug Res.* 2003; 53(4): 266-271.

34. Ra CS, Jung BY, Park G. The Fungicidal Benzothiazole Methoxyacrylates: Synthesis, Conformational Analysis and Fungicidal Activity. *Heterocycles*. 2004; 62(1): 793-802.
35. Lochart A, Ye L, Judd DB et al. Evidence for the Presence of Three Distinct Binding Sites for the Thioflavin T Class of Alzheimer's Disease PET Imaging Agents on β -Amyloid Peptide Fibrils *J. Biol. Chem.* 2005; 280(9): 7677-7684.
36. Bradshaw TD, Stevens MFG. The Discovery of the Potent and Selective Antitumour Agent 2-(4-Amino-3-methylphenyl) benzothiazole (DF 203) and Related Compounds. *Curr. Med. Chem.* 2001; 8(2): 203-210.
37. Wenzel SE, Kamada AK. Zileuton: the first 5-lipoxygenase inhibitor for the treatment of asthma. *Ann. Pharmacother.* 1996; 30(7-8): 858-864.
38. Jutchinson I, Jennings SA, Vishnuvajjala BR, Westwell AD, Stevens MFG. Antitumor Benzothiazoles. 16.¹ Synthesis and Pharmaceutical Properties of Antitumor 2-(4-Aminophenyl)benzothiazole Amino Acid Prodrugs. *J. Med. Chem.* 2002; 45(3): 744-747.
39. Green AG, Perkin AG. Polythiosulphonic acids of *p*-diamines. *J. Chem. Soc. Trans.* 1903; 83: 1201-1212.
40. Landquist JK. Diaminobenzobisthiazoles and Related Compounds. *J. Chem. Soc.(C)*. 1967; 2212-2220.
41. Kiprianov AI, Mikhailenko FA. Synthesis and structures of isomeric benzobisthiazoles. *Khim. Geterotsikl. Soedin.* 1967; 3: 270; original ref.: *Chem. Heterocycl. Compds.* 1969; 3(4): 205-209.
42. Solar SL, Cox RJ, Clecak NJ, Ettinger R. Synthesis and proof of structure of 2,6-diamino-benzo[1,2-d:4,5-d']bisthiazole. *J. Org. Chem.* 1968; 33(5): 2132-2133.
43. Cullen E, Becker R, Freter K et al. Bis Basic Substituted Diaminobenzobisthiazoles as Potential Antiarthritic Agents. *J. Med. Chem.* 1992; 35(2): 350-361.
44. Hegershoff H. *Ber.* 1903; 36, 3121.
45. Ambati NB, Anand V, Hanumantha P. A Facile Synthesis of 2-N(Methyl Amino) Benzothiazoles. *Synth. Commun.* 1997; 27(9): 1487-1493.
46. Edge SRH. Benzobisthiazoles. *J. Chem. Soc. Trans.* 1922; 121: 772-775.
47. Edge SRH. Benzobisthiazoles. II. *J. Chem. Soc. Trans.* 1923; 123: 153-156.
48. Edge SRH. Benzobisthiazoles. III. *J. Chem. Soc. Trans.* 1923; 123: 1011-1014.
49. Finzi C, Grandolini G. *Gazz. Chim. Ital.* 1959; 89: 2543; *Chem. Abstr.* 1961; 55: 27841.

50. Neunhoeffer O, Barnikow G. Über Benzo-bis-thiazole. Chem. Ber.1962; 95(3): 616-619.
51. Barnikow G, Kunzek H, Hoffmann M. Zur Kenntnis der HUGERSHOFF-Reaktion. I. Die Cyclisierung von m-Phenylen-bis-thioharnstoff. J. Prakt. Chem.1965; 27(5-6): 271-275.
52. Grandolini G. Gazz. Chim. Ital.1960; 90: 1221; Chem. Abstr. 1962; 56: 31395.
53. Jacobson P. Ber. Dtsch. Chem. Ges.1886; 19: 1067.
54. Finzi C, Grandolini G. Gazz. Chim. Ital.1959; 89: 2555; Chem. Abstr. 1961; 55: 27842.
55. Grandolini G, Ricci A, Martani A, Mezzetti T. Thiocyanation of amino-2-methylbenzothiazoles. J. Het. Chem.1966; 3(3): 299-301.
56. Grandolini G, Ricci A, Martani A, Monache FD. Nuclear magnetic resonance of benzobisheterocyclic compounds. J. Het. Chem.1966; 3(3): 302-310.
57. Grandolini G, Martani A, Fravolini A. Ann. Chim.1968; 58: 1248; Chem. Abstr. 1969; 70: 57724.
58. Bogert MT, Taylor WS. Researches on thiazoles. XVII. An investigation of the connection between constitution and color in the thioflavine group. Collect. Czech. Chem. Commun.1931; 3: 480-498.
59. Stephens FF, Wibberley DG. Some potential trypanocidal and antibacterial compounds in the heterocyclic series. Part I. Benzothiazoles. J. Chem. Soc.1950; 3336-3340.
60. Grandolini G, Martani A, Ricci A, Bellavait C. Ann. Chim.1968; 58: 91; Chem. Abstr. 1968; 69: 19099.
61. Chakrabarty M, Mukherji A, Karmakar S et al. An expedient synthesis of novel 2-substituted thiazolo [4,5-f] isoquinolines/quinolines and benzo [1,2-d:4,3-d'] bisthiazoles and their potential as inhibitors of COX-1 and COX-2. ARKIVOC. 2010; (xi): 265-290.
62. Mike JF, Inteman JJ, Ellern A, Malika Jeffries-EL M. Facile Synthesis of 2,6-Disubstituted Benzobisthiazoles: Functional Monomers for the Design of Organic Semiconductors. J. Org. Chem. 2010; 75(2): 495–497.
63. Kimura K, Meurer D, Hutzler RF et al. Synthesis and Characterization of Thermotropic Poly(benzobisthiazole)s. Macromolecules. 1994; 27(6): 1303-1306.
64. Harrington PJ, Lodewijk E. Twenty Years of Naproxen Technology. Org. Process Res. Dev. 1997; 1(1): 72-76.
65. Ziakas GN, Rekka EA, Gavalas AM, Eleftheriou PT, Kourounakis PN. New analogues of butylated hydroxytoluene as anti-inflammatory and antioxidant agents. Bioorg. Med. Chem. 2006; 14(16): 5616-5624.

66. Geronikaki AA, Lagunin AA, Hadjipavlou-Litina DI et al. Computer-Aided Discovery of Anti-Inflammatory Thiazolidinones with Dual Cyclooxygenase/Lipoxygenase Inhibition. *J. Med. Chem.* 2008; 51(6): 1601-1609.
67. Li B, Pai R, Aiello D. Optimization of a novel potent and selective bacterial DNA helicase inhibitor scaffold from a high throughput screening hit. *Bioorganic & Medicinal Chemistry Letters.* 2013; 23(12): 3481-3486.
68. Prak K, Kriston-Vizi J, Edith Chan AW et al. Benzobisthiazoles Represent a Novel Scaffold for Kinase Inhibitors of CLK Family Members. *Biochemistry.* 2016; 55 (3): 608-617.
69. Čibová A, Magdolen P, Fülöpová A et al. Pushpull molecular structures based on angular benzobisthiazolium acceptor. *Synt. Photophys. Prop. Their. Stud. Tetrahedron.* 2015; 71: 315–323.
70. Ahmed E, Kim FS, Xin H, Jenekhe SA. Benzobisthiazole-Thiophene Copolymer semiconductors: Synthesis, Enhanced Stability, Field-Effect Transistors, and Efficient Solar Cells. *Macromolecules.* 2009; 42(22): 8615–8618.
71. Alam MM, Jenekhe SA. Polybenzobisazoles Are Efficient Electron Transport Materials for Improving the Performance and Stability of Polymer Light-Emitting Diodes. *Chem. Mater.* 2002; 14(11): 4775-4780.
72. Osaheni JA, Jenekhe SA. New red light-emitting conjugated rigid-rod polymer: poly(benzobisthiazole-1,4-phenylenebisvinylene). *Macromolecules.* 1993; 26(17): 4726-4728.
73. Jenekhe SA, Osaheni JA, Meth JS, Vanherzeele H. Nonlinear optical properties of poly (p-phenylenebenzobisoxazole). *Chem. Mater.* 1992; 4(3): 683-687.
74. Osaheni JA, Jenekhe SA. Synthesis and processing of heterocyclic polymers as electronic, optoelectronic, and nonlinear optical materials. 1. New conjugated rigid-rod benzobisthiazole polymers. *Chem. Mater.* 1992; 4(6): 1282-1290.

75. Dotrong M, Mehta R, Balchin GA et al. Synthesis, processing, and third-order nonlinear optical properties of benzobisthiazole polymers containing thiophene moieties. *J. Polym. Sci. Part A*. 1993; 31(3): 723-729.
76. Osaheni JA, Jenekhe SA, Perlstein J. Efficient photogeneration of charge carriers in a layered photoreceptor using a conjugated polymer. *J. Appl. Phys. Lett.* 1994; 64(23): 3112.
77. Osaheni JA, Jenekhe SA, Perlstein J. Photogeneration of charge carriers in bilayer assemblies of conjugated rigid-rod polymers. *J. Phys. Chem.* 1994; 98(48): 12727–12736.
78. Jenekhe SA, Yi S. Highly Photoconductive Nanocomposites of Metallophthalocyanines and Conjugated Polymers. *Adv. Mater.* 2000; 12(17): 1274-1278.
79. Zhang X, Jenekhe SA. Electroluminescence of Multicomponent Conjugated Polymers. 1. Roles of Polymer/Polymer Interfaces in Emission Enhancement and Voltage-Tunable Multicolor Emission in Semiconducting Polymer/Polymer Heterojunctions. *Macromolecules*. 2000; 33(6): 2069-2082.
80. Kawamoto T, Takeda K, Nishiwaki M, Aridomi T, Konno T. Square-Planar $N_2S_2Ni^{II}$ Complexes with an Extended π -Conjugated System. *Inorg. Chem.* 2007; 46(10): 4239-4247.
81. Hu XB, Yu DM. Hyperbranched Polybenzobisthiazole with High Thermal Stability, Good Organosolubility, and Interesting Optical Performance. *Macromol. Chem. Phys.* 2012; 213(7): 738-746.
82. Kang JG, Kim HJ, Jeong YK et al. Optical and Conformational Studies on Benzobisthiazole Derivatives. *J. Phys. Chem. B*. 2010; 114(11): 3791-3798.
83. Babel A, Jenekhe SA. *n*-Channel Field-Effect Transistors from Blends of Conjugated Polymers. *J. Phys. Chem. B*. 2002; 106(24): 6129-6132.
84. Wang L, Wada T, Yuba T et al. Nonlinear optical anisotropy and molecular orientational distribution in poly(p-phenylene benzobisthiazole) Langmuir-Blodgett films. *J. Appl. Phys.* 1996; 79(12): 9321-9325.
85. Lee JW, Wang CS. Processing of optical quality rigid-rod polymer thin films. *Polymer*. 1994; 35(17): 3673–3678.
86. Roberts MF, Jenekhe SA, Cameron A, Mcmillan M, Perlstein J. Lewis Acid Coordination Complexes of Polymers. 2. Computational Modeling of Single-Chain and Aggregate Structures of Rigid-Rod Poly(p-phenylenebenzobisthiazole). *Chem. Mater.* 1994; 6(5): 658-670.

87. Bai SJ, Spry RJ, Zelmon DE, Ramabadran U, Jackson J. Optical anisotropy of polymeric films measured by waveguide propagation mode determination. *J. Polym. Sci. Part B Polym. Phys.* 1992; 30(13): 1507-1514.
 88. Vanherzeele H, Meth JS, Jenekhe SA, Roberts MF. Third-order nonlinear optical properties of thin films of poly(*p*-phenylene benzobisthiazole) and its molecular composites with polyamides. *Appl. Phys. Lett.* 1991; 58: 663-665.
 89. Reinhardt BA, Unroe MR, Evers RC. Third-order optical nonlinearities of model compounds containing benzobisthiazole, benzobisoxazole, and benzbisimidazole units. *Chem. Mater.* 1991; 3(5): 864-871.
 90. Osaheni JA, Jenekhe SA, Perlstein J. Efficient photogeneration of charge carriers in a layered photoreceptor using a π -conjugated polymer. *Appl. Phys. Lett.* 1994; 64(23): 3112-3114.
 91. Ahmed E, Subramaniyan, S, Kim FS, Xin H, Jenekhe SA. Benzobisthiazole-based donor-acceptor copolymer semiconductors for photovoltaic cells and highly stable field-effect transistors. *Macromolecules.* 2011; 44(18): 7207-7219.
 92. McEntee GJ, Vilela F, Skabara PJ, Anthopoulos TD, Labram JG, Tierney S, Harrington RW, Clegg W. Self-assembly and charge transport properties of a benzobisthiazole end-capped with dihexyl thienothiophene units. *J. Mater. Chem.* 2011; 21(7): 2091–2097.
 93. Bhuwarka A, Mike JF, He M, Intemann JJ, Nelson T, Ewan MD, Rogers RA, Lin Z, Jeffries-El M. Quaterthiophene-Benzobisazole Copolymers for Photovoltaic Cells: Effect of Heteroatom Placement and Substitution on the Optical and Electronic Properties. *Macromolecules.* 2011; 44(24): 9611-9617.
-