

## *International Journal of Scientific Research and Reviews*

### **A Study on the Occurrence of Species Diversity *Pediastrum* in Selected Temple Tanks in Chennai, Tamil Nadu, India.**

**D. Karthikeyan, S.Kumaresan and B.S. Balakumar\***

Post Graduate and Research Department of Plant Biology & Plant Biotechnology, Ramakrishna Mission Vivekananda College, Chennai - 600004, India.

\*E-mail: [bsbviveka@gmail.com](mailto:bsbviveka@gmail.com)

#### **ABSTRACT**

A periodical survey on the occurrence of the green algae *Pediastrum* in the temple tanks of Arulmigu Sri. Parthasarathy swamy Temple, Thiruvallikeni, Chennai, and Arulmigu Sri Kapaleeswarar Temple, Mylapore, Chennai, India, was done during the period October 2016 to September 2017. A maximum of six species of *Pediastrum* along with physico-chemical parameters such as: colour, odour, turbidity, electrical conductivity, calcium, sodium, chloride, fluoride levels and TDS of water samples were recorded. The study showed interesting variations in physico-chemical parameters and species availability during the study period.

**KEY WORD:** *Pediastrum*, Temple tank, Microalgae, Physico-chemical parameters.

#### **\*Corresponding author**

**Dr. B.S.Balakumar**

Associate Professor

Post Graduate and Research

Department of Plant Biology & Plant Biotechnology,

Ramakrishna Mission Vivekananda College,

Chennai - 600004, India.

E-mail: [bsbviveka@gmail.com](mailto:bsbviveka@gmail.com), Mobile No: 9444667500

## INTRODUCTION

Green algae are aquatic plants that act as pioneer photosynthetic organisms or producers, in an aquatic ecosystem. The genus *Pediastrum* Mayen, belongs to class Chlorophyceae and order Sphaeropleales. It is a free floating, coenobial, green algae occurring commonly in natural freshwater lentic environments like ponds, lakes, reservoirs.<sup>1</sup> However their occurrence in brackish and salty waters is rare.<sup>2</sup> The outer layer of cell wall of *Pediastrum* is composed of sporopollenin combined with oxides of silicon which makes them highly resistant to decay. Therefore, they remain preserved well in lake sediments as fossil record for palynological studies.<sup>3</sup> After the monograph of Philipose, many more species of *Pediastrum* have been added to the Indian algal flora. Eleven species were reported from the Indian region.<sup>4</sup>

Presence of pollution tolerant algae like *Melosira*, *Oscillatoria*, *Pediastrum* and *Scenedesmus* has been considered as indicative of enriched waters, thus providing evidence of pollution of water.<sup>5</sup>

*Scenedesmus* was the diverse genera with seven species followed by *Pediastrum* with five species and *Schroederia* and *Phacus* represented by three species each. Fifteen pollution tolerant algal species were identified from this pond during the period of study.<sup>6</sup>

The genus *Pediastrum* was reported four species in birds visited ponds of kannyakumari<sup>7</sup> and the recorded eight *Pediastrum* species in kerala temple tanks.<sup>8</sup>

## MATERIALS AND METHODS

Algal and water samples collected from *Arulmigu Sri. Parthasarathy swamy* Temple, Thiruvallikeni, Chennai, india, tank located at 13° 3' 14.0724" N ; 80° 16' 37.2396" E and *Arulmigu Sri Kapaleeswarar* Temple , Mylapore, Chennai, India, located at 13° 2' 0.1176" N ; 80° 16' 7.41" E. The temple tank water samples were collected, during October 2016 to September 2017. The samples were collected in clean plastic containers, brought to the laboratory and divided into two parts, one part was preserved in 4% formalin, the water samples were analysed according to American Public Health Standards.<sup>9</sup>

Observations for micro-algal presence were made using light microscope unit. Taxonomical descriptions of taxa were determined on the basis of morphological features such as outline of cells, number of lobes and processi, depth of incisions in marginal cells, sculpture of cell wall. Cell dimensions are described as: width x length. In the case of marginal cells the length of a cell includes length of lobes and processi. All observations were made using oil immersion objective lens (10x X 100x Magnification). Identification of the microalgae was done using monograph on Chlorococcales.<sup>4</sup>

## RESULT

**Table 1: List of *Pediastrum* species identified from samples collected from Parthasarathy swamy Temple & Kapaleeswarar Temple.**

S.No	Name of the <i>Pediastrum</i> species	Parthasarathy swamy Temple tank			Sri Kapaleeswarar Temple tank		
		M	W	S	M	W	S
1	<i>Pediastrum simplex</i> Meyen	+	-	-	+	+	+
2	<i>Pediastrum simplex</i> var. <i>duodenarium</i> (J.W.Bailey) Rabenhors	-	-	-	+	+	-
3	<i>Pediastrum ovatum</i> (Ehrenberg) A.Braun	-	-	-	+	-	-
4	<i>Pediastrum duplex</i> var. <i>genuinum</i> (A.Braun) Hansgirg	-	-	+	-	-	-
5	<i>Pediastrum duplex</i> var. <i>clathratum</i> (A.Braun) Lagerheim	-	+	+	-	-	-
6	<i>Pediastrum duplex</i> var. <i>reticulatum</i> Lagerheim	+	+	+	-	-	-
7	<i>Pediastrum duplex</i> var. <i>gracillimum</i> West & G.S.West	+	-	+	-	-	-
8	<i>Pediastrum tetras</i> var. <i>tetraodon</i> (Corda) Hansgirg	+	+	+	+	+	+
9	<i>Pediastrum tetras</i> var. <i>apiculatum</i> Fritsch	+	+	+	+	+	-

M-Monsoon, W-Winter, S-Summer

Present +, Absent -

**Table:2 Physico-chemical parameters of water samples collected from Parthasarathy swamy Temple Tank & Kapaleeswarar Temple Tank**

S.No	Parameters-mg/L	Parthasarathy swamy Temple tank			Kapaleeswarar Temple tank		
		Monsoon	Winter	Summer	Monsoon	Winter	Summer
1	Appearance	Slightly greenish	Slightly greenish	Slightly greenish	Slightly greenish	Slightly greenish	Slightly greenish
2	Odour	Agreeable	Agreeable	Agreeable	Agreeable	Agreeable	Agreeable
3	Turbidity	Turbid	Turbid	Turbid	Turbid	Turbid	Turbid
4	Electrical Conductivity (micro mho/cm)	244	230	225	285	260	275
5	pH at 25°C	7.05	7.4	7.5	7.01	7	8.72
6	Alkalinity total	92	90	90	60	50	50
7	Total hardness	74	40	50	50	50	80
8	Calcium	26	19	20	16	14	17
9	Magnesium	2	1	2	2	2	1
10	Sodium	24	20	18	34	25	18
11	Potassium	2	2	1	3	2	1
12	Iron	0	0	0	0	0	1
13	Manganese	0	0	0	0	0	0
14	Free ammonia	5	2	3	3	0.5	0.5
15	Nitrite	0.1	0.1	0	0.5	0	0.1
16	Nitrate	1	1	0	1	1	0
17	Chloride	28	30	50	57	60	130
18	Fluoride	0.5	0.5	0.5	0.2	1	1.5
19	Sulphate	0	0	0	0	0	0
20	Phosphate	0.5	0.2	0.3	0.5	0.5	0.2
21	Residual chlorine	0.2	0.2	0.2	2.1	0.2	0.2
22	Total TDS	171	192	228	200	240	312

## Species Identification

(Based on the monograph on chlorococcales by M.T. Philipose, 1967)

Phylum: *Chlorophyta*

Subphylum: *Tetraphytina*

Class: *Chlorophyceae*

Order: *Sphaeropleales*

Family: *Hydrodictyaceae*

Subfamily: *Hydrodictyoideae*

Genus: *Pediastrum* Meyen,

*Pediastrum* colonies are free-floating, disc-or stellate -shaped, generally of 4-8-16-32 more polygonal multinucleate cells arranged in a single layer one cell thick. Colony with or without perforations. (Marginal cells are with one, two or four processes, rarely without such processes). Cell wall is smooth, granulate or with reticulate ridges. (Chloroplast, single parietal, fills the adult cells with one or more pyrenoids). Asexual reproduction by zoospores which escape into a vesicle and orient themselves in the place of cells, rarely by 1-2 hypnospores are produced from each cell.

Eleven species are known from Indian region.

## Taxonomic Description

### 1. *Pediastrum simplex* Meyen (Pl.2 Fig.1a,b)

M.T. Philipose 1967, p.113. fig.36a,36b.

Colonies are circular to oval, of 4-8-16-32 or more cells. Inner side of marginal cells nearly straight or outer side produced into a gradually tapering process with, sides concave. Inner cells are polygonal. Cells are in contact with adjacent ones and are usually without intercellular spaces. When present, intercellular spaces are very small. Cell wall smooth or punctuate to granulate. Cells are (7-) 8-13  $\mu\text{m}$  broad and (15-) 19-26 (-30)  $\mu\text{m}$  long.

### 2. *Pediastrum simplex* var. *duodenarium* (J.W.Bailey) Rabenhors ( Pl.2 Fig.2a,2b )

M.T. Philipose 1967 p.115. fig.36d,e

It differs from the type in having large intercellular spaces or a single central space with the cell arranged in a ring at the periphery. (Inner face of marginal cells are concave and outer face is prolonged into a single delicately tapering process). Sides of marginal cells also concave or nearly straight. Interior cells similar to marginal cells but with shorter processes. Cell wall is smooth or finely punctuate. Colonies are composed of 4-8-16-32-64-128 (usually 8-16-32) cells. Cells being 8-24  $\mu\text{m}$  broad, and 10-45  $\mu\text{m}$  long; if 16-celled colonies are 125  $\mu\text{m}$  in diameter.

### 3. *Pediastrum ovatum* (Ehrenberg) A.Braun (Pl.2 Fig.3 )

M.T. Philipose 1967, p.115. fig.37d.

Colonies, usually 4-8-16-(rarely 32-) celled, with the cells are arranged in a ring round a central space or with one or more interior cells and a number of marginal cells, perforate or almost imperforate, the perforations being small. Cells are plumper than in *P.simplex* var. *duodenarium* with the outer sides of periphery and often central cells convex. Cell wall is smooth or ornamented. Four-celled colonies up to 60  $\mu$ , 8-celled colonies up to 30  $\mu$  and 16-celled colonies up to 100  $\mu$  in diameter, each cell measuring 8.5-19  $\mu$ m broad and 14-37  $\mu$ m long.

4. *Pediastrum duplex* Meyen var. *genuinum* ( A. Braun) Hansgirg ( Pl.2 Fig.4 )

M.T. Philipose 1967, p.123. fig.43d.

Colonies are 4-8-16-32 celled with fairly large intercellular spaces. Marginal cells are with stout processes which are straight or slightly curved, with cell membrane smooth or punctate. Cells measure 6-18  $\mu$ m , perforations up to 6  $\mu$ m and colonies upto 45-65  $\mu$ m in diameter

5. *Pediastrum duplex* Meyen var. *clathratum* ( A. Braun) Lagerheim (Pl.2 Fig.5 )

M.T. Philipose 1967, p.123. fig.43f.

Cells have deeply emarginate sides and large intercellular spaces in *Pediastrum duplex*. Colonies are 3-64 celled, cells being 9-25  $\mu$ m in diameter. Sixteen –celled colonies measure up to 90  $\mu$ m in diameter.

6. *Pediastrum duplex* Meyen var. *reticulatum* Lagerheim (Pl.2 Fig.6 )

M.T. Philipose 1967, p.124. fig.43g.

Cells have ( 10-20 (-40)  $\mu$ m in diameter) more or less H-shaped with sides of processes of marginal cells nearly parallel. Intercellular spaces are large and oval. Colonies 8-16 celled are 58-70  $\mu$ m in diameter.

7. *Pediastrum duplex* Meyen var. *gracillimum* W. Et G. S. West ( Pl.2 Fig.7 )

M.T. Philipose 1967, p.124. fig.43i.

Colonies with very large intercellular spaces have very narrow cells, as broad as or narrower than the processes. Body of marginal cells are curved outwards with two long processes having emarginated apices. Inner cells also similar to marginal cells but with shorter processes. Cells are 10-18.5 (-22)  $\mu$ m broad and 12-25 (-32)  $\mu$ m long with Perforations 4-16  $\mu$ m in diameter. Sixteen celled colonies measure 65-140  $\mu$ m in diameter.

8. *Pediastrum tetras* (Ehr.) Ralfs var. *tetraodon* ( Corda ) Hansgirg (Pl.2 Fig.8a,b)

M.T. Philipose 1967, p.129. fig.45d,g.


Colonies 4-8-16 celled. Incision of cells deep with the lobes adjacent to the incision of the marginal cells very pronounced. Cells 8-18  $\mu$ m in diameter.

9. *Pediastrum tetras* (Ehr.) Ralfs var. *apiculatum* Fritsch (Pl.2 Fig.9)



M.T. Philipose 1967, p.130. fig.45h.

The colonies from Azhicode had processes with apical nodular thickenings as seen in *P. tetras* var. *apiculatum* Fritsch. The colony was 24.6 µm in diameter with cells 14-17.6 µm in diameter. Owing to insufficiency of material it could not be decided whether the alga belonged to this type.



**Plate: 1.**Photomicrographs of *pediatrion* species recorded in the water samples collected from Parthasarathy swamy Temple and Kapaleeswarar Temple.

1a,b. *Pediatrion simplex* Meyen 2a,b. *Pediatrion simplex* var. *duodenarium* (J.W.Bailey) Rabenhors 3. *Pediatrion ovatum* (Ehrenberg) A.Braun 4. *Pediatrion duplex* var. *genuinum* (A.Braun) Hansgirg 5. *Pediatrion duplex* var. *clathratum* (A.Braun) Lagerheim 6. *Pediatrion duplex* var. *reticulatum* Lagerheim 7. *Pediatrion duplex* var. *gracillimum* West & G.S.West 8a,b. *Pediatrion tetras* var. *tetraodon* (Corda) Hansgirg 9. *Pediatrion tetras* var. *apiculatum* Fritsch.

## DISCUSSION

Presence of algae like *Melosira*, *Oscillatoria*, *Pandorina*, *Pediatrion*, *Closterium*, *Navicula*, *Microcystis* and *Scenedesmus* has been considered indicative of nutrients - enriched water providing evidence of pollution.<sup>10</sup> High population by *Cosmarium* *Scenedesmus*, *Pediatrion*, and *Pleurotaenium* species was observed during the period of August to October.<sup>11</sup>

In the present study, the water samples collected from temple tanks of *Arulmigu Sri. Parthasarathy swamy* Temple tank, Thiruvallikeni, Chennai, and *Arulmigu Sri Kapaleeswarar* Temple tank, Mylapore, Chennai, India, for a year, recorded a total of nine *Pediastrum* species. The maximum number of *Pediastrum* species was recorded in *Parthasarathy swamy* Temple tank during Summer. The species *Pediastrum tetras* var. *tetraodon* (Corda) Hansgirg was found to be present in both the temple tanks in all the seasons studied. The physico-chemical parameters of the water sample of *Kapaleeswarar* temple tank recorded maximum levels in Total TDS and Chloride during the summer season, contributing to the observed diversity of the alga. The pollution of the study area is due to increased drain of pollutants into the tanks by visiting devotees in summer.

## ACKNOWLEDGEMENT

The authors gratefully acknowledge the help and facilities extended by the Secretary, RKM. Vivekananda College, Mylapore, Chennai – 600 004 for carrying out the study.

## REFERENCES

1. Meyen, F.I.F. Beobachtungen ueber einige niedere algenformen. Nova Acta Phy. Med. Acad. Cass. Leop. Carol. Nat. Curr. 1829; 14(2): 768-78.
2. Parra, O.O., Revision der gattung Pediastrum Meyen (*Chlorophyta*). *Bibl. Phycol* 1979; 48: 1-186.
3. Komárek, J. and V. Jankovská., Review of the green algae genus Pediastrum; Implication for pollen-analytical research. *Bibl. Phycol.*, Cramer J. Berlin-Stuttgart, 2001; 108.
4. Philipose MT., *Chlorococcales*. Indian Council of Agricultural Research, New Delhi. India, 1967; 1–365.
5. Tessy, P.P. and Sreekumar, R., A report on the pollution algae from the Thrissur Kol wetlands (part of Vembanad-Kol, Ramsar site), Kerala. *Nat. Environ. Polln. Tech*, 2008; 7: 311–314.
6. Tessy Paul and Anu P.K., Algal diversity of Guruvayur Temple Pond, Thrissur District, Kerala. *International Journal of Advanced Life Sciences (IJALS)*. 2016; 9(3): 302-306.
7. Regini Balasingh, S. Jeeva & W. Vincy., Species Composition of Phytoplankton in the Two Migratory Birds Visited Ponds of Kanyakumari District. *International Journal of Botany*. 2015; 5 (5):1-8
8. Arulmurugan, Subramani Nagarajand Narayanswamy Anand., Biodiversity of Fresh Water Algae From Temple Tanks of Kerala. *Recent Research in Science And Technology*. 2010; 2 : 58-71.



9. APHA., Standard methods of Examination of Waste water, AWAA, WPCF Washington, DC, USA. 1985; 20 :1-1268.
10. Biswas, B.K. and Konar, S.K., Impact of waste disposal on plankton abundance and diversity in the River Ganga at Hatdidah (Bihar). Poll. Res. 2000; 19:633-640.
11. Amit Kumar and Radha Sahu., Diversity of Algae (Cholorophyceae) in Paddy Fields of Lalgutwa Area, Ranchi, Jharkhand. *Journal of Applied Pharmaceutical Science*. 2012; 2: 092-095.